


Direct (e^+, e^-) pair creation by muon

Direct (e^+ , e^-) pair creation by muon

Creation of a (e^+ , e^-) pair by virtual photon in the Coulomb field of the nucleus (for momentum conservation).



It is one of the most important processes of muon interaction.

At TeV muon energies, pair creation cross section exceeds those of other muon interaction processes in a wide region of energy transfers :

$$100 \text{ MeV} \leq \epsilon \leq 0.1 E_\mu$$

Average energy loss for pair production **increases linearly** with muon energy, and in TeV region this process contributes over 50 % to the total energy loss rate.

energy transfers

Main contribution to the **total cross section** is given by transferred energies:


$$5\text{MeV} \leq \epsilon \leq 0.01 E_\mu$$


The contribution to **average muon energy loss** is determined mostly by region:

$$10^{-3} E_\mu \leq \epsilon \leq 0.1 E_\mu$$

Thus, to adequately describe the number of pairs produced, average energy loss and stochastic energy loss distribution one needs to reproduce with a sufficient accuracy the differential cross section behaviour in a wide range of energy transfers:

$$5\text{MeV} \leq \epsilon \leq 0.1 E_\mu$$





differential cross section

The differential cross section is given by Kokoulin et al. [Koko71].

It includes :

- screening of the field of the nucleus
- correction for finite nuclear size
- contribution from the atomic electrons [Keln97]
- ...

See [Koko71] for a complete discussion.

differential cross section

The differential cross section per atom can be written as :

$$\frac{d\sigma}{d\epsilon} = \frac{4 \alpha^2 r_e^2}{3\pi} \frac{1-v}{\epsilon} [Z(Z + \zeta)] F(Z, E, \epsilon) \quad (1)$$

with

$$F(Z, E, \epsilon) = \int_0^{\rho_{max}} \left[\Phi_e(v, \rho) + \left(\frac{m_e}{m_\mu} \right)^2 \Phi_\mu(v, \rho) \right] d\rho \quad (2)$$

where

$$\epsilon = \epsilon^+ + \epsilon^- = \text{total energy of the created pair;}$$

$$v = \epsilon/E$$

$$\rho = (\epsilon^+ - \epsilon^-)/\epsilon = \text{asymmetry coefficient;}$$

The functions Φ_e , Φ_μ , ζ can be found in [Koko00].

limits

$$\epsilon_{min} = 4m_e c^2$$

$$\epsilon_{max} = E - \frac{3\sqrt{e}}{4} m_\mu c^2 Z^{1/3}$$

$$\rho_{min} = 0$$

$$\rho_{max} = \left[1 - \frac{6(m_\mu c^2)^2}{E(E - \epsilon)} \right] \sqrt{1 - \frac{\epsilon_{min}}{\epsilon}}$$

Energetic pairs and truncated energy loss rate

One may wish to take into account separately the high-energy pairs emitted **above a given threshold** ϵ_{cut} (miss detection, explicit simulation ...).

Those pairs must be **excluded** from the mean energy loss count.

$$\left. -\frac{dE}{dx} \right]_{\epsilon < \epsilon_{cut}} = n_{at} \int_{\epsilon_{min}}^{\epsilon_{cut}} \epsilon \frac{d\sigma}{d\epsilon} d\epsilon$$

n_{at} is the number of atoms per volume.

Then, the truncated total cross-section for emitting 'hard' pairs is:


$$\sigma(E, \epsilon_{cut} \leq \epsilon \leq \epsilon_{max}) = \int_{\epsilon_{cut}}^{\epsilon_{max}} \frac{d\sigma}{d\epsilon} d\epsilon$$

The muon deflection angle is of the order of:

$$\theta = \frac{mc^2}{E}$$


Above ~ 1000 TeV the LPM suppression mechanism may have an effect.

number of interactions per cm in Iron. (cut 100 MeV)



muon kinetic energy (GeV)



1 TeV muon in 10 meter of Fe (field 5 tesla).
direct pair creation only




10 meter of Fe : muons 100 GeV, 1 TeV, 5 TeV.

left : brems only


right : brems + direct pair creation



energy deposited by high energy muons in a bloc of Iron



energy spectrum of 1 TeV muons after 3 m of iron



References

- [Lohm85] W.Lohmann,R.Kopp,R.Voss, Cern Report 85-03 (1985)
- [Koko69] R.P.Kokoulin and A.A.Petrukhin, Proc. 11th Intern. Conf. on Cosmic Rays, Budapest, 1969 [Acta Phys. Acad. Sci. Hung.,**29**, Suppl.4,p.277, 1970].
- [Koko71] R.P.Kokoulin and A.A.Petrukhin, Proc. 12th Int. Conf. on Cosmic Rays, Hobart, 1971, **vol.6**, p.2436.
- [Keln97] S.R.Kelner, Phys. Atomic Nuclei, **61** (1998) 448.
- [Koko00] S.R.Kelner, R.P.Kokoulin, A.Rybin. Geant4 Physics Reference Manual, Cern (2000)
- [PDG] D.E. Groom et al. Particle Data Group . Rev. of Particle Properties. Eur. Phys. J. C15,1 (2000) <http://pdg.lbl.gov/>